


4TH EDITION

Biological Safety

Principles and
Practices



Edited by
Diane O. Fleming and
Debra L. Hunt

Copyright © 2006

ASM Press
American Society for Microbiology
1752 N Street, N.W.
Washington, DC 20036-2804

Library of Congress Cataloging-in-Publication Data

Biological safety: principles and practices / editors, Diane O. Fleming, Debra L. Hunt.—4th ed.
p. ; cm.

Includes bibliographical references and index.

ISBN-13: 978-1-55581-339-0

ISBN-10: 1-55581-339-9

1. Microbiological laboratories—Safety measures. 2. Biological laboratories—Safety measures.
I. Fleming, Diane O. II. Hunt, Debra Long.

[DNLM: 1. Accidents, Occupational—prevention & control. 2. Laboratories—standards.
3. Safety. WA 485 B615 2006]

QR64.7.L33 2006

570'.289—dc22

2006019406

All Rights Reserved

Printed in the United States of America

10 9 8 7 6 5 4 3 2 1

Address editorial correspondence to: ASM Press, 1752 N St., N.W., Washington, DC 20036-2904, U.S.A.

Send orders to: ASM Press, P.O. Box 605, Herndon, VA 20172, U.S.A.

Phone: 800-546-2416; 703-661-1593

Fax: 703-661-1501

Email: Books@asmusa.org

Online: estore.asm.org

Laboratory, Growth Chamber, and Greenhouse Microbial Safety: Plant Pathogens and Plant-Associated Microorganisms of Significance to Human Health

ANNE K. VIDAVER, SUE A. TOLIN, AND PATRICIA LAMBRECHT

3

Several years ago, interkingdom pathogenic specificity was rarely recognized (Starr and Chatterjee, 1972). We now know that an increasing number of the organisms, and occasionally even the same strains of an organism, can infect both plants and humans. Commonalities in gene sequences and function among pathogens of plants, animals, and humans are no longer surprising (Cao et al., 2001; Alfano and Collmer, 2004; Govan and Vandamme, 1998; Govan, et al., 1996; Gorbalenya, et al., 1989; Rodriguez-Villalobos et al., 2002; Tan, 2002). In addition, some plant-associated microorganisms which may prevent plant disease nevertheless can cause human allergies or disease. Such biocontrol agents are regulated commercially by the U.S. Environmental Protection Agency for environmental risks but not for risks to humans. Hence, it is prudent to assess safety issues with respect to human exposure to plant-associated microorganisms, including viruses, for laboratory and confined uses. Keeping with the purpose of this book, our focus is on risk characterization and mitigation of worker exposure during culturing, inoculation of plants, and diagnosis of plant pathogens known to affect human health.

For bacteria, commonalities among plant, animal, or human pathogens are most evident in type III secretion pathways (Alfano and Collmer, 2004; Hueck, 1998) and virulence factors in pseudomonads (Govan and Vandamme, 1998; Rahme et al.,

1995; Tan, 2002). In most cases, virulence or pathogenicity factors in common still await discovery. In fungi, commonalities are at the structural, morphological, biochemical, and genetic levels (Hall et al., 1999; Procop and Roberts, 1998). In viruses, gene sequences may be highly homologous and similarities in genomic functionality are known (Gorbalenya et al., 1989; Hohn and Fütterer, 1997; Toh et al., 1983).

The emergence or reemergence of human diseases caused by microorganisms is due to many factors (Vidaver, 1996). In humans, for example, both *Burkholderia cepacia* and *Pseudomonas aeruginosa* have become important pathogens in cystic fibrosis; both are also an infrequent cause of infection in non-cystic fibrosis patients (Holmes et al., 1999; Vikram et al., 1999). The emergence of newly identified fungal pathogens and the reemergence of previously uncommon fungal diseases of humans (mycoses) are attributed to an increase in the number of susceptible individuals, such as bone marrow and organ transplant recipients, cancer patients being treated with chemotherapy, critically ill persons, very-low-birth-weight infants, and persons with certain other infections, notably human immunodeficiency virus (Dixon et al., 1996). Clinically relevant mycoses may occur in healthy, immunocompetent individuals as well (Pontón et al., 2000; see also chapter 8 on mycotic diseases).

The many texts and manuals that deal with methods of working with plant pathogens do not provide any cautions or statements with respect to potential risk to human health (Blanchard and Tater, 1981; Burgess and Liddell, 1983; Dhingra and Sinclair, 1995; Fahy and Persley, 1983; Hampton et al., 1990; Hickey, 1986; Kahn and Mathur, 1999; Klement et al., 1990; Lelliot and Stead, 1997; Razin and Tully, 1983; Rechcigl and Rechcigl, 1997; Saettler et al., 1989; Schaad, 2001; Schneck, 1982; Singleton et al., 1992; Tuite, 1988; CDC/NIH, 1999; NIH, 2002; U.S. Department of Agriculture [USDA], 1992), except for a one-page statement in a recent handbook (Ritchie, 2002). Nor in medical texts is there any caution about the potential for occupational exposure to plant pathogens or plant-associated organisms, other than as allergens (Horner et al., 1995).

There are increasing numbers of plant-pathogenic microbial organisms associated with human diseases or maladies (Table 1). Of the more than 300 reported species of fungi isolated from humans with infectious systemic diseases (Taylor et al., 2001), at least 40 are known plant pathogens. Of these, all but two are in the phylum Ascomycota, popularly known as ascomycetes. Multiple species in the genera *Alternaria*, *Aspergillus*, *Bipolaris*, *Colletotrichum*, *Curvularia*, and *Fusarium* have been implicated in human disease. Ten additional genera are represented by one species each. This suggests specificity in evolution of pathogenicity as well as susceptibility in humans, which may simply be the ability of humans to meet the nutritional and asexual reproductive capacity or other characteristics of these fungi. Many plant-pathogenic fungi are associated with mycotoxicoses, but these are not considered in Table 1 because entry is via consumption of contaminated foods and unlikely to occur in a laboratory setting (Bennett and Klich, 2003).

Of the more than 500 species of bacteria isolated from human infections (Taylor et al., 2001), 12 are also known as plant pathogens or as biocontrol agents associated with plants. Reports of human infections for 22 bacterial species are listed in Table 1. Of these species, only six are gram positive, with three being *Bacillus* spp. The rest are gram negative and include multiple species in the genera *Agrobacterium*, *Burkholderia*, *Enterobacter*, *Klebsiella*, *Pseudomonas*, and *Serratia*, and single species of *Pantoea*, *Stenotrophomonas*, and *Xanthomonas*. Homologues of bacterial genes coding for virulence or pathogenicity factors in common between crossover pathogens are becoming better known (Table 2) (Alfano and Collmer, 2001; Tan, 2002). For access to the literature, as well as to journals covering plant diseases or associations, the reader is referred to common texts in plant pathology

(Agrios, 2005; Lucas and Dickinson, 1998). Only a few direct references combine plant and human microbial associations.

No report to date implicates plant viruses in human disease. Anecdotally, plant virologists have been known to produce antibodies to viruses with which they have worked, and have shown allergic sensitivities following the inhalation of viral aerosols. However, it is unlikely that a plant virus could infect and become viremic in humans. Specific proteins on the outer surface of animal and human viruses bind to cellular receptors, proteins on the outer surface of cells, inducing penetration, translocation of viral components to intracellular sites, and replication (Mettenleiter, 2002). In contrast, plant viruses have no known cellular receptor-binding proteins and require a mechanically induced or insect-induced wound in a host cell to facilitate penetration. Theoretically, a needle stick from a hollow-bore needle attached to a syringe containing a plant pathogen could also provide a mode of entry into the human host.

Given the small size of viral genomes, and similarity in functions, it is not surprising to see many closely related sequences between plant and human viruses (Table 2). There has been one suggestion that a plant nanovirus had recombined with picorna-like viral RNA to form circoviruses infecting vertebrates (Gibbs and Weiller, 1999). The host switch recombinational event is postulated to have occurred in a vertebrate when it was exposed to sap from a nanovirus-infected plant. Thus, plant viruses may pose a risk in the event that there is an open wound or percutaneous injury through which a virus may enter. The end result, however, is likely to be the creation of a new virus which may or may not cause disease in the infected host. Dimitrov (2004) and Baranowski et al. (2001) reviewed virus entry and the evolution of cell recognition by viruses and suggested that minimal changes in viral genomes may trigger changes in receptor usage for virus entry; neither review considered plant viruses.

RISK ASSESSMENT AND BLs

As the NIH guidelines note (NIH, 2002), risk assessment is ultimately a subjective process. By the standard that agents are not associated with disease in healthy adult humans, almost all the plant pathogens and plant-associated microorganisms are in NIH risk group 1, for which biosafety level 1 (BL1) is usually recommended. However, as noted in Table 1, a few plant pathogens should be viewed as more problematic in that some strains of some species may infect both immunocompromised hosts and, more rarely,

TABLE 1 Continued

| Taxon ^a | Plant disease/association | Human disease |
|---|--|--|
| <i>Stenotrophomonas maltophilia</i> | Plant-associated and plant pathogen (Suckstorff and Berg, 2003) | Bacteremia and respiratory tract infections (Denton and Kerr, 1998; Goss et al., 2004) |
| <i>Xanthomonas campestris</i> pv. <i>campestris</i> | Black rot of crucifers and wilt and blight stump rot of broccoli, cabbage, cauliflower, brussel sprouts, kale, mustard, radish, rutabaga, sunflower, stock, and turnip (Agrios, 2005; Westcott, 2001) | Bacteremia (Li et al., 1990) |
| Fungi^b | | |
| Phylum Ascomycota <i>Alternaria alternata</i> | Wide host range. Causes leaf spots, blights, damping off, and stem and fruit rots (Westcott, 2001). Black mold of tomato, leaf spot and ear and root rot of maize, blight of foliage and pod of pea, seedling foliage blight of sugarcane (Rott and Comstock, 2000), black point on wheat, dark flecks on <i>Dendrobium</i> , fruit spot on papaya, and leaf spot on garden bean | Phaeohyphomycosis (Duffill and Coley, 1993), mycotic keratitis, cutaneous (Ono et al., 2004) and visceral infections, and osteomyelitis |
| <i>Alternaria tenuissima</i> | Strawberry fruit rot (Howard and Albregts, 1973), leaf spot of broad bean (Honda et al., 2001), leaf spot of high bush blueberry (Milholland, 1995), leaf spot of <i>Amaranthus hybridus</i> (Blodgett and Swart, 2002), and leaf spot of papaya | Phaeohyphomycosis (Romano et al., 1996; Romano et al., 1997), sinusitis, and ulcerated cutaneous and visceral infections (Rossman et al., 1996) |
| <i>Aspergillus candidus</i> | Decay of apple fruits (Thind et al., 1976), root rots | Cerebral aspergillosis (Linares et al., 1971), cutaneous aspergillosis, endocarditis, endophthalmitis, hepatosplenic aspergillosis, meningitis, myocarditis, onychomycosis, osteomyelitis, otomycosis, pulmonary aspergillosis (Krysinska-Traczyk and Dutkiewicz, 2000), and sinusitis |
| <i>Aspergillus flavus</i> | Pathogen and saprophyte; has many hosts and causes such diseases as ear and kernel rot of maize (St. Leger et al., 2000; Smart et al., 1990), yellow mold of peanut (Pitt et al., 1991), and boll rot on cotton (Brown et al., 1992) | Systemic aspergillosis (Yamada et al., 1998) and endocarditis (Rao and Saha, 2000) |
| | Biopesticide: <i>A. flavus</i> strain AF36 (Arizona Cotton Research and Protection Council, Phoenix, Ariz.), a non-toxin-producing strain registered (EPA ^c) on cotton fields in Texas and Arizona for control of strains of <i>A. flavus</i> which produce aflatoxin; <i>A. flavus</i> strain NRRL 21882, registered for use in peanut crops to control aflatoxin-producing strains of <i>A. flavus</i> (Circle One Global, Inc., Shellman, Ga.) | |

(TABLE 1 continued)

TABLE 1 Taxa of pathogens and saprophytes of plants associated with human disease or maladies

| Taxon ^a | Plant disease/association | Human disease |
|---|---|--|
| Bacteria | | |
| Gram positive | | |
| <i>Bacillus megaterium</i> | White blotch of wheat and bacterial wetwood of poplar and elm (Murdoch and Campana, 1983) | Oral mucosal inflammation (Rubinstein and Pedersen, 2002) |
| <i>Bacillus circulans</i> | Date palm disease (Leary and Chun, 1989) | Peritonitis (Berry et al., 2004), serious nongastrointestinal infections in animals, and diarrheal enterotoxin production in human cells (Rowan et al., 2003; Deva and Narayan, 1989) |
| <i>Bacillus pumilus</i> | Bacterial blotch of immature Balady peach (Saleh et al., 1997); registered biocontrol agent, strain GB34 Yield Shield; (Gustafson, Plano, Tex.) for control of soilborne fungal pathogens causing root disease in soybean | Oral mucosal inflammation (Rubinstein and Pedersen, 2002; Suominen et al., 1999) |
| <i>Clostridium butyricum</i> | Wetwood of poplar (Schink et al., 1981) and disease of hornbeam (Gvozdiak et al., 1976) | Necrotizing enterocolitis in babies (Howard et al., 1977) |
| <i>Clostridium histolyticum</i> | Plant associated | Gas gangrene (myonecrosis) and necrotic lesions (Brazier et al., 2004) |
| <i>Rathayibacter toxicus</i> (syn. <i>Clavibacter toxicus</i>) | Gummosis of cereals (Riley and Ophel, 1992) | Death of livestock associated with consumption of <i>Rathayibacter</i> -infected annual ryegrass (Riley and Ophel, 1992); human disease speculative (Edgar, 2004) |
| Gram negative | | |
| <i>Agrobacterium radiobacter</i> (syn. <i>Rhizobium radiobacter</i>) | Plant-associated bacteria, rhizosphere; registered biocontrol agent for crown gall, strain K84 (Galltrol A; AgBioChem Inc., Orinda, Calif.) in fruit, nut, and ornamental nursery stock and strain K1026 (Nogall; Bio-Care Technology Pty Ltd., Somersby, New South Wales, Australia) for control of crown gall on fruit and nut trees, caneberries, roses, and other ornamentals | Opportunist pathogen (Edmond, 1993). Bacterial endophthalmitis (Miller et al., 1996); bacteremia (Southern, 1996), endocarditis (Plotkin, 1980), peritonitis (Melgosa Hijosa et al., 1997), and urinary tract infections (Namdari et al., 2003; Dunne et al., 1993) |
| <i>Agrobacterium tumefaciens</i> (syn. <i>Rhizobium tumefaciens</i>) | Agent of crown gall with wide host range (Moore and Warren, 1979; Westcott, 2001) | Peritonitis (Ramirez et al., 1992; Alnor et al., 1994), bacteremias (Southern, 1996), and urinary tract infection (Hulse et al., 1993) |
| <i>Burkholderia cepacia</i> | Sour skin of onion (Burkholder, 1950; Yohalem and Lorbeer, 1997) and cavity disease of <i>Agaricus bitorquis</i> (Gill and Cole, 1992; Alameda and Migrucci, 1998); phytoremediation (Glick, 2004) and endophyte (Hinton and Bacon, 1995) | Bacteremia (Woods et al., 2004); pulmonary complex (De Boeck et al., 2004; Courtney et al., 2004); serious respiratory pathogen in cystic fibrosis patients (Wigley and Burton, 1999; Govan et al., 1996); bacteremia, cardiac cirrhosis and cellulitis (Lau et al., 1999); and endophthalmitis (Pathengay et al., 2004) |

(TABLE 1 continued)

TABLE 1 Continued

| Taxon ^a | Plant disease/association | Human disease |
|---|---|--|
| <i>Burkholderia gladioli</i> | Slippery skin of onion (Kishun and Swarup, 1981); decay of <i>Gladiolus</i> spp., <i>Iris</i> spp., and rice; leaf spot and blight of <i>Asplenium nidus</i> (Chase et al., 1984); and bacterial disease of <i>Dendrobium</i> sp. orchid (Chuenchitt et al., 1983) | Bacteremia (Shin et al., 1997), pneumonia (Ross et al., 1995), and cervical adenitis (Graves et al., 1997) |
| <i>Enterobacter cloacae</i> | Wetwood on elm, internal decay of onion (Bishop and Davis, 1990), and rhizome rot of edible ginger (Nishijima et al. 2004); biocontrol agent (Punja, 1997; Wilson et al., 1987; Watanabe et al., 2000) | Septicemia and respiratory tract infections (Jochimsen et al., 1998) and gas gangrene (Fata et al., 1996) |
| <i>Erwinia persicina</i> (syn. <i>Erwinia nulandii</i>) | Necrosis in fruits, vegetables (Hao et al., 1990), necrosis of bean pods and seeds (Brenner et al., 1994) | Urinary tract infection (O'Hara et al., 1998) |
| <i>Klebsiella pneumoniae</i> | Endophyte; many plant hosts, including maize (Dong et al., 2003) | Pneumonia (Prince et al., 1997), bacteremia (Kang et al., 2004), and meningitis (Tang and Chen, 1994) |
| <i>Klebsiella variicola</i> | Plant-associated on banana, rice, sugarcane, and maize (Rosenblueth et al., 2004) | Bacteremia and urinary tract infection (Rosenblueth et al., 2004) |
| <i>Pantoea agglomerans</i> (syn. <i>Enterobacter agglomerans</i> , <i>Erwinia herbicola</i>) | Pathogen of <i>Wisteria</i> and onion; wetwood of elm; black flesh of pineapple and grapefruit; spot disease and frost damage on corn, soy, and clover; and disease of millet (Frederickson et al., 1997). Saprophyte | Nosocomial/opportunistic infections (Bennett et al., 1995) and septic arthritis (Kratz et al., 2003) |
| <i>Pseudomonas aeruginosa</i> | Onion rot (Cothier et al., 1976) and <i>Arabidopsis</i> rot | Burn wound infections and pneumonia (Rahme et al., 1995; Johansen et al., 1998; Vikram et al., 1999) and meningitis, bacteremia, and sepsis (Torii et al., 2003) |
| <i>Pseudomonas fluorescens</i> | Registered biocontrol for <i>Erwinia amylovora</i> (Johnson and Stockwell, 1998) on apple, cherry, almond, peach, and pear (Blight Ban A506; Frost Technology Corporation, Burr Ridge Ill.); frost protection on fruit crops, almond, tomato, and potato to reduce frost-forming bacteria on leaves and blossoms (Frostban; Frost Technology Corporation) | Bacteremia (Hsueh et al., 1998) |
| <i>Pseudomonas putida</i> | Plant saprophyte with potential for application in biological control of plant pathogens, bioremediation, and production of bioplastics (Nelson et al., 2002) | Nosocomial infections (Lombardi et al., 2002); meningitis (Ghosh et al., 2000); and bacteremia, pneumonia, and sepsis (Torii et al., 2003) |
| <i>Serratia ficaria</i> | Plant associated (biological cycle) (Grimont et al., 1979) | Organ infections (Anahory et al., 1998) and endophthalmitis, gall bladder empyema, and septicemia (Badenoch et al., 2002) |
| <i>Serratia marcescens</i> | Alfalfa crown and root rot (Lukezic et al., 1982), cucurbit yellow vine disease (Bruton et al., 2003), and endophytic colonization of rice (Gyaneshwar et al., 2001) | Respiratory tract infections, urinary tract infections, and bacteremia (Ostrowsky et al., 2002) and conjunctivitis, endocarditis, meningitis, and wound infections (Su et al., 2003) |

(TABLE 1 continued)

TABLE 1 Continued

| Taxon ^a | Plant disease/association | Human disease |
|---------------------------------------|---|--|
| <i>Aspergillus glaucus</i> | Corn ear and kernel rot (Nyvall, 1979) | Cerebral, cutaneous, hepatosplenic, and pulmonary aspergillosis; endocarditis; endophthalmitis; meningitis; myocarditis; onychomycosis; osteomyelitis; otomycosis; and sinusitis (O'Shaughnessy et al., 2004) |
| <i>Aspergillus niger</i> | Black mold of peanut (Nyvall, 1979) and onion (Tanaka and Nonaka, 1977) and maize ear rot (Nyvall, 1979) | Aspergilloma (fruiting body in tissue), otomycoses (Mishra et al., 2004), and pulmonary aspergillosis (Yamaguchi et al., 1992) |
| <i>Aspergillus oryzae</i> | Saprophyte and mycotoxin producer (Geiser et al., 2000) | Necrotizing scleritis (Stenson et al., 1982) and bronchopulmonary aspergillosis (Akiyama et al., 1987) |
| <i>Aureobasidium pullulans</i> | Russet of apple fruit (Matteson Heidenreich et al., 1997) and d'Anjou pear (Spotts and Cervantes, 2002) and stem break and browning of flax | Various opportunistic mycoses, pulmonary mycoses, scleritis (Gupta et al., 2001), and phaeohyphomycosis (Kaczmarek et al., 1986) |
| <i>Bipolaris australiensis</i> | Leaf spot and crown and root rot of turfgrass, (http://www.apsnet.org/online/common/comment/turfgrass.asp) | Phaeohyphomycosis, allergic and chronic sinusitis, keratitis, endophthalmitis (Chalet et al., 1986), endocarditis, osteomyelitis, meningitis, encephalitis, peritonitis, and pulmonary infection (Flanagan and Bryceson, 1997) |
| <i>Bipolaris hawaiiensis</i> | Leaf and culm lesions on Callides Rhodesgrass (Sonoda, 1991) and Bermuda grass disease (Pratt, 2001) | Endophthalmitis, phaeohyphomycotic orbitopathy, sinusitis, and granulomatous encephalitis (Morton et al., 1986) |
| <i>Bipolaris spicifera</i> | Leaf spot of cotton (http://www.apsnet.org/online/common/names/cotton.asp) | Phaeohyphomycosis (McGinnis, et al., 1992), fungal endarteritis (Ogden et al., 1992), meningitis (Latham, 2000), and peritonitis (Bava et al., 2003) |
| <i>Chaetomium globosum</i> | Disease in tomato (Gerald et al., 1980) and infection of barley roots | Cerebral phaeohyphomycosis (Anandi et al., 1989) and onychomycosis (Hattori et al., 2000) |
| <i>Cladosporium oxysporum</i> | Leaf spots and blights of many plants and leaf spot of pepper (Hammouda, 1992) | Phaeohyphomycosis (Romano et al., 1999) and pneumonia (Yeghen et al., 1996) |
| <i>Colletotrichum coccodes</i> | Black dot of tomato (Dillard and Cobb, 1997) and potato (Andrion et al., 1998) | Phaeohyphomycosis (O'Quinn et al., 2001) |
| <i>Colletotrichum gloeosporioides</i> | Anthrax on many fruits and plantation crops (Westcott, 2001), including anthracnose of papaya leaves (Dickman and Alvarez, 1983) and avocado, poplar, aspen, and cottonwood shoot blight, fruit rot on apple and berries of coffee, and dieback of citrus | Keratitis (Yamamoto et al., 2001) and phaeohyphomycosis (O'Quinn et al., 2001) |
| <i>Coniothyrium fuckelii</i> | Stem blight, dieback, and canker of <i>Rosa</i> spp. and strawberry (<i>Fragaria</i>) and black root rot and cane blight of <i>Rubus</i> spp. (Heimann and Boone, 1983) | Mycotic keratitis (Laverde et al., 1973) and liver infection (Kiehn et al., 1987) |
| <i>Curvularia brachyspora</i> | Leaf spot disease of <i>Rosa</i> spp. (Kore and Bhide, 1976) | Necrotizing cutaneous infection (Torda and Jones, 1997) and mycotic keratitis (Marcus et al., 1992) |
| <i>Curvularia clavata</i> | Leaf spot of maize (Mandokhot and Chaudhary, 1972) | Invasive sinusitis and cerebritis (Ebright et al., 1999) and human skin infection (Gugnani et al., 1990) |

(TABLE 1 continued)

TABLE 1 Continued

| Taxon ^a | Plant disease/association | Human disease |
|-----------------------------------|---|---|
| <i>Curvularia geniculata</i> | Banana leaf spot (Meredith, 1963) and melting out of turfgrasses (Westcott, 1990) | Mycotic keratitis (Georg, 1964) and maduromycotic mycetomas in animals (Bridges, 1957) |
| <i>Curvularia lunata</i> | Leaf rot of rice (Lakshmanan, 1992), leaf spot of bentgrass (Muchovej and Couch, 1987), melting out of turfgrasses (Westcott, 2001), leaf spot of maize (Ito et al., 1979), and leaf spot of cotton (Gour and Dube, 1975) | Cerebral phaeohyphomycosis (Carter and Boudreaux, 2004), systemic cutaneous infection (Tessari et al., 2003), and allergic fungal rhinosinusitis (Taj-Aldeen et al., 2004) |
| <i>Curvularia pallescens</i> | Leaf spot of sugarcane (Rao et al., 1992), leaf spot and ear rot of maize (Lal and Tripathi, 1977), brown spot of asparagus, and leaf spot of rubber | Phaeohyphomycosis (Agrawal and Singh, 1995) |
| <i>Curvularia senegalensis</i> | Seedling foliage blight on sugarcane (Rott and Comstock, 2000) and leaf spot of maize and other hosts (Yang, 1973) | Mycotic keratitis (Guarro et al., 1999) |
| <i>Cylindrocarpon lichenicola</i> | Postharvest fruit invasion and corm rot of <i>Colocasia esculenta</i> (taro) (Usharani and Ramarao, 1981) | Disseminated infection (Rodriguez-Villalobos et al., 2003) and keratomycosis (Mangiaterra et al., 2001) |
| <i>Drechslera biseptata</i> | <i>Drechslera</i> leaf spot of turfgrasses and black point in wheat grains (Fischl et al., 1993); mycotoxin producer (Leach and Tulloch, 1972) | Brain abscess (Mycology Online, 2004) |
| <i>Fusarium chlamydosporum</i> | Root rot and wilt of <i>Coleus forskohlii</i> (Boby and Bagyaraj, 2003) and blight of kangaroo paw (<i>Anigozanthos</i> spp.) (Satou et al., 2001) | Invasive infection (Segal et al., 1998) |
| <i>Fusarium dimerum</i> | One of several agents of fig endosepsis (Michailides et al., 1996) | Disseminated infection (Austen et al., 2001), endocarditis (Camin et al., 1999), and eye infection (Vismer et al., 2002) |
| <i>Fusarium incarnatum</i> | Walnut canker (Seta et al., 2004) and aster wilt | Mycotic keratitis (Naiker and Odhav, 2004); black gill disease of shrimp (Khoja et al., 2004) |
| <i>Fusarium moniliforme</i> | Ear, root, and stalk rot and seedling blight of maize (Westcott, 2001); sugarcane wilt complex; and pseudostem heart rot of banana (Jones and Lomeiro, 2000); wide host range | Human fusariosis, local and systemic (Dignani and Anaissie, 2004) |
| <i>Fusarium oxysporum</i> | Wilts and blights on a wide range of vegetable and plantation crops, ornamentals, small grains (Bottalico and Perrone, 2002), and turfgrasses, including potato, sugarcane, bean, cowpea, and <i>Musa</i> spp. (Raabe et al., 1981), and corm and root rots (Lucas and Dickinson, 1998) | Disseminated fusariosis (Sander et al., 1998), skin and nail infection (Romano et al., 1998), pneumonia (Rodriguez-Villalobos et al., 2002), and onychomycosis (Godoy et al., 2004) |
| <i>Fusarium proliferatum</i> | Leaf, sheath, stem spots, damping off, and flower spots on <i>Dendrobium</i> and <i>Cattleya</i> orchid; head blight in wheat and other small-grain cereals (Bottalico and Perrone, 2002); and wilt and dieback of date palm (Abdalla et al., 2000) | Disseminated infection in immunosuppressed individuals (Summerbell et al., 1988) and suppurative thrombophlebitis (Murray et al., 2003) |

(TABLE 1 continued)

TABLE 1 Continued

| Taxon ^a | Plant disease/association | Human disease |
|---------------------------------|---|--|
| <i>Fusarium solani</i> | Yellows, fruit rots, seedling rots, root rots, and damping off on a wide range of hosts; fungal root rot of banana (Jones and Stover, 2000); and stem canker of sweet potato, black walnut, and poinsettia (Westcott, 2001) | Invasive furiosis (Repiso, et al., 1996; Bushelman et al., 1995) and onychomycosis (Godoy et al., 2004) |
| <i>Lasiodiplodia theobromae</i> | Fruit and stem rot of papaya (Dantas et al., 2003); canker of dogwood (Mullen et al., 1991); kumquat dieback (Ko et al., 2004); black kernel rot of maize; crown, finger, stalk, and peduncle rot of banana (Abeywickrama et al., 2004); and collar rot of peanut (Phipps and Porter, 1998) | Subcutaneous abscess (Maslen et al., 1996), ophthalmic mycoses (Thomas, 2003), onychomycosis, and phaeohyphomycosis |
| <i>Lecythophora hoffmannii</i> | Soft rots and decay of the surface layers of natural and preservative-treated timber (Bugos et al., 1988) | Chronic sinusitis (Marriott et al., 1997) |
| <i>Paecilomyces variotii</i> | Dieback and canker of pistachio (Ashkan et al., 1997) | Pneumonia (Byrd et al., 1992), central nervous system infection (Kantarcioglu et al., 2003), and peritonitis (Wright et al., 2003) |
| <i>Phoma eupyrena</i> | Blight of fir and pine species (Kliejunas et al., 1985) | Cutaneous lesions (Bakerspigel et al., 1981). |
| Phylum Zygomycota | | |
| <i>Mucor circinelloides</i> | Fruit rot of <i>Luffa acutangula</i> (Singh et al., 1974) and mucor rot of mango (http://www.ismpminet.org/resources/common/names/mango.asp) | Zygomycosis (Chandra and Woodgyer, 2002) and gangrenous mucormycosis (Boyd et al., 2003) |
| <i>Rhizopus oryzae</i> | Fruit rots of pineapple, mango, and carrot (http://www.ismpminet.org/resources/common/names) | Pulmonary zygomycosis (Eisen and Robson, 2004) |
| <i>Rhizopus stolonifer</i> | Pre- and postharvest soft rots of many fruits, vegetables, and crops; sunflower head rot (Yang et al. 1979); and seedling blight on lupine (Westcott, 2001) | Zygomycosis (Gonzalez et al., 1996) |

^aUnidentified or inadequately identified species of *Microbacterium* and *Streptomyces* have been reported for clinical infections (Funke et al., 1997; Carey et al., 2001), and both identified and inadequately identified species have been associated with plants (Kaku, 2004; Zinniel et al., 2002; Westcott, 2001).

^bNot included are zoonotic fungi, those from animals that can cause infections in people, fungal biocontrol agents, and commercial fungi used in brewing or baking. Members of these classes can cause human clinical disease. Toxigenic fungi and mycotoxins are not addressed.

^cEPA, Environmental Protection Agency.

immunocompetent hosts. Sources of pathogens for microbial infection and contamination of plants include infected or infested seed, wind-driven inoculum, contaminated harvest machinery and containers, irrigation water, and postharvest handling (Scholthof, 2003). Humans can be exposed to microorganisms via any of these environmental sources. Exposure of humans is likely to increase during laboratory procedures of pathogen culturing and isolation. The airborne spores produced by fungi can initiate mycoses if inhaled by humans. By-products of

plant pathogens in food, such as mycotoxins, can also cause illness (Bennett and Klich, 2003). Allergic reactions to plant pathogens and products such as toxins, while known to us, are not well documented (Hall et al., 1999). Allergens from some plant-pathogenic fungi, primarily *Alternaria* and *Fusarium* species, were included in a review by Horner et al. (1995).

With the prevalent paradigm of specificity among plant and human pathogens, it is not surprising that the literature is sparse on comparative connections among taxonomic groups of pathogens of different

TABLE 2 Selected fungal and viral gene homologues coding for virulence or pathogenicity factors in plant and human diseases

| Gene(s) | Plant pathogen/host | Organisms known to cause disease or disease/malady |
|---|---|---|
| SP1; codes for extracellular protein (Hall et al., 1999) | <i>Stagonospora (Septoria) nodorum</i> ; wheat blotch | <i>Coccidioides immitis</i> , <i>Aspergillus fumigatus</i> , respiratory allergenic proteins |
| Polymerases, RNase H, RNA-binding ZN finger, protease, primer-binding site (Hohn and Fütterer, 1997; Nault, 1997; Toh et al., 1983) | Cauliflower mosaic virus; many crop hosts; pararetroviruses | Retroviruses, hepadnaviruses, poliovirus |
| Helicase, protease, RNA-dependent RNA polymerase | Cherry rasp leaf virus (James and Upton, 2002; Thompson et al., 2004); Satsuma dwarf virus (Karasev et al., 2001) | Calici- and picornaviruses not known in humans, Equine rhinitis B virus, foot-and-mouth disease virus (Thompson et al., 2004) |

hosts (Govan and Vandamme, 1998; Rahme et al., 1995; Hueck, 1998). Few studies have examined the capacity of microorganisms to cause disease in both plants and animals (Rahme et al., 1995; Wigley and Burton, 1999; Govan and Vandamme, 1998; Tan, 2002). It has long been recognized that certain plant-infecting viruses are able to propagate in insect vectors and that viruses in the same family (e.g., *Reoviridae* and *Rhabdoviridae*) are pathogens of vertebrates (Nault, 1997).

RISK POTENTIAL FOR LABORATORY PERSONNEL

As mentioned previously, healthy adults are not normally at risk of being infected by plant-associated microorganisms, but allergic reactions may occur. Good laboratory practices and some specific suggestions for dealing with plant-associated microorganisms are found in the methods manuals previously cited and the NIH guidelines. Immunocompromised adults, e.g., transplant recipients and those with immunodeficiencies (genetic or microbial), and persons with allergic sensitivities should take particular care in handling plant pathogens and microorganisms associated with plants. Large-scale cultures, aerosol-generating procedures, the use of needles and syringes, and direct contact with skin wounds are examples of activities that may increase the risk of exposure and infection.

CONTAINMENT

Samples of plant material obtained for isolation of pathogens or biocontrol agents are protected from contamination by using aseptic techniques, including

surface sterilization of seeds, leaves, stems, and roots. In isolation procedures, materials may be ground or sliced to obtain the putative organism directly or after concentration in centrifuge tubes; buffers are usually added to such materials to obtain suitable suspensions and to provide optimal pH and ionic composition for stabilizing the structure of the pathogen. Some pathogens are obligate parasites or simply cannot be cultured; such organisms are not currently known to present an infectious risk to humans, but they may be allergenic. After microorganisms are isolated, experimental procedures frequently involve the generation of aerosols, e.g., by flaming, low- or high-pressure spraying, or inoculating plants with various mechanical devices. Plants may need to be wounded prior to or during inoculation with such materials as silicon carbide or Carborundum, which itself is a corneal and respiratory irritant.

Air-purifying particulate filter respirators that are effective against intake of particles 2 μm or larger from the surrounding air should be considered for all personnel and recommended for at-risk groups who may be exposed to potentially infectious or allergenic aerosols. The organisms known to have been associated with a disease condition in humans, and thus of particular concern, are listed in Table 1. The N95 NIOSH (National Institute for Occupational Safety and Health) series of filters are recommended. They are easy to wear, disposable, and of modest cost. In scientific supply catalogs, these filters have NIOSH approval numbers with the prefix TC-21C. Surgical masks are not suitable, since they fail performance criteria for protection against airborne contaminants. (Further information on respiratory protection may be found in chapter 17.)

The recommended containment level for all plant-associated microorganisms is BL1 or BL2 conditions in the laboratory or growth chamber, and BL1-P or BL2-P conditions in the greenhouse, as delineated in the NIH guidelines (NIH, 2002). These principles of containment are applicable to both wild-type and recombinant organisms and are design based to protect laboratory workers. When BL3-P conditions are required by regulators, they are for minimizing escape of the pathogen and for protection of the environment, not people.

Guidelines for using microorganisms in field work or natural ecosystems are provided by the USDA (1992) and can be found in the primary literature on plant pathology. The texts and methods manuals mentioned in the introduction may be consulted for organismal isolation, survival, growth, decontamination, and plant inoculation. Each organism-plant interaction is unique and requires special conditions of plant susceptibility, such as plant age, tissue specificity, temperature, humidity, and photoperiod, for achieving an infection which mimics natural conditions. Specific insect vectors may be necessary to inoculate plants with some viruses that cannot be mechanically transmitted. For several purposes, tobacco (*Nicotiana tabacum*) and the weed *Arabidopsis thaliana* are considered model plants for testing putative plant pathogens, akin to mouse models for human pathogens. There is only one documented case of endangerment of field populations of plants known to have occurred as a result of the use of plant pathogens in contained facilities (McKeen, 1989). Nevertheless, the USDA's Plant Protection Act regulations (section 7CFR330) require that unless a plant pathogen has been isolated locally, permits for its use must be obtained and containment conditions specified therein must be followed. These conditions are aimed at preventing damage to plants through environmental dissemination, rather than at protecting the human worker.

DISPOSAL

In the laboratory or greenhouse, autoclaving cultures and pathogen-infected material, or otherwise rendering them biologically inactive, is routine. In gardens and experimental fields, as well as commercial areas, timely chemical treatments for a few bacteria, many fungi, and some insect vectors or wild hosts decreases the inoculum. Chemicals are not available or cost-effective for controlling many plant pathogens, and biocontrol agents are few. Other management practices to decrease the inoculum, and thus decrease exposure, are crop rotation, planting of resistant varieties (where available), planting date, and plowing

under infected or infested plant material. These practices decrease inoculum by the process of competition with other microorganisms in the soil, where many plant pathogens are poor survivors. Composting can also be effective.

MOVEMENT OF PLANT PATHOGENS

If the isolated microorganism is known to be a plant pathogen (e.g., see Agrios, 2005), irrespective of risk (no official risk groups are delineated for risk of a pathogen to plants or for risk to humans), a permit is required from the USDA-APHIS to move the agent from one location, state, or country to another. Packaging, storage, and transportation of plant pathogens are also under the aegis of APHIS rules and regulations (USDA, 2001), as well as the Department of Commerce. However, suspected or unknown pathogens can be sent by ordinary mail to laboratories for diagnostic purposes, as is done for human clinical specimens. (See also chapter 21 on shipping.)

CONCLUDING REMARKS

Greater caution should be exercised by those working with plant pathogens and biocontrol agents than has routinely been taken. Specifically, aerosol generation in plant and culture experimentation should be minimized or carefully controlled. In addition to concern about the etiologic agent, multiple antibiotic resistance is exhibited by some of these microorganisms, such as *B. cepacia* (Wigley and Burton, 1999) and *Stenotrophomonas maltophilia* (Denton and Kerr, 1998). For fungi, mortality from invasive infections is much greater than for bacterial pathogens (Engelhard, 1998), due to the limited treatment options available. Immunocompromised adults and persons with allergic sensitivities or open wounds should take particular care in handling plant pathogens and microorganisms associated with plants. Development of rapid and reliable methods for identifying viral, bacterial, and fungal pathogens will make diagnosis easier. New therapeutic approaches for treating invasive bacterial and fungal infections of humans, as well as virus-associated allergies, are future medical challenges.

References

- Abdalla, M. Y., A. Al-Rokibah, A. Moretti, and G. Mulè. 2000. Pathogenicity of toxigenic *Fusarium proliferatum* from date palm in Saudi Arabia. *Plant Dis.* **84**: 321-324.
- Abeywickrama, A. S., R. Dayananda, S. W. Wijeratnam, and L. Arambewela. 2004. Fungal pathogens associated

- with banana fruit in Sri Lanka, and their treatment with essential oils. *Mycopathologia* 157:91–97.
- Agrawal, A., and S. M. Singh. 1995. Two cases of cutaneous phaeohyphomycosis caused by *Curvularia pallescens*. *Mycoses* 38:301–303.
- Agrios, G. N. 2005. *Plant Pathology*, 5th ed. Elsevier Academic Press, Burlington, Mass.
- Akiyama, K., H. Takizawa, M. Suzuki, S. Miyachi, M. Ichinohe, and Y. Yanagihara. 1987. Allergic bronchopulmonary aspergillosis due to *Aspergillus oryzae*. *Chest* 91:285–286.
- Alameda, M., and J. S. Mignucci. 1998. *Burkholderia cepacia*, causal agent of bacterial blotch of oyster mushroom. *J. Agric. Univ. P. R.* 82:109–110.
- Alfano, J. R., and A. Collmer. 2001. Mechanisms of bacterial pathogenesis in plants: familiar foes in a foreign kingdom, p. 179–226. In E. A. Groisman (ed.), *Principles of Bacterial Pathogenesis*. Academic Press, San Diego, Calif.
- Alfano, J. R., and A. Collmer. 2004. Type III secretion system effector proteins: double agents in bacterial disease and plant defense. *Annu. Rev. Phytopathol.* 42:385–414.
- Alnor, D., N. Frimodt-Moller, F. Espersen, and W. Frederiksen. 1994. Infections with the unusual human pathogens *Agrobacterium* species and *Ochrobactrum anthropi*. *Clin. Infect. Dis.* 18:914–920.
- Anahory, T., H. Darbas, O. Ongaro, H. Jean-Pierre, and P. Mion. 1998. *Serratia ficaria*: a misidentified or unidentified rare cause of human infections in fig tree culture zones. *J. Clin. Microbiol.* 36:3266–3272.
- Anandi, V., T. J. John, A. Walter, J. C. Shastry, M. K. Lalitha, A. A. Padhye, L. Ajello, and F. W. Chandler. 1989. Cerebral phaeohyphomycosis caused by *Chaetomium globosum* in a renal transplant recipient. *J. Clin. Microbiol.* 27:2226–2229.
- Andrison, D., J. M. Lucas, C. Guérin, and B. Jouan. 1998. Colonization of roots, stolons, tubers and stems of various potato (*Solanum tuberosum*) cultivars by the black dot fungus *Colletotrichum coccodes*. *Plant Pathol.* 47:440–445.
- Ashkan, S. M., D. Abusaidi, and D. Ershad. 1997. Etiological study of dieback and canker of pistachio nut tree in Rafsangan. *Iran. J. Plant Pathol.* 33:15–26.
- Austen, B., H. McCarthy, B. Wilkins, A. Smith, and A. Duncombe. 2001. Fatal disseminated *Fusarium* infection in acute lymphoblastic leukaemia in complete remission. *J. Clin. Pathol.* 54:488–490.
- Bakerspigel, A., D. Lowe, and A. Rostos. 1981. The isolation of *Phoma eupyrena* from a human lesion. *Arch. Dermatol.* 117:362–363.
- Badenoch, P. R., A. L. Thom, and D. J. Coster. 2002. *Serratia ficaria* endophthalmitis. *J. Clin. Microbiol.* 40:1563–1564.
- Baranowski, E., C. M. Ruiz-Jarabo, and E. Domingo. 2001. Evolution of cell recognition by viruses. *Science* 292:1102–1105.
- Bava, A. J., A. Fayad, C. Cespedes, and M. Sandoval. 2003. Fungal peritonitis caused by *Bipolaris spicifera*. *Med. Mycol.* 41:529–531.
- Bennett, J. W., and M. Klich. 2003. Mycotoxins. *Clin. Microbiol. Rev.* 16:497–516.
- Bennett, S. N., M. M. McNeil, L. A. Bland, M. J. Arduino, M. E. Villarino, D. M. Perrotta, D. R. Burwen, S. F. Welbel, D. A. Pegues, L. Stroud, P. S. Zeitz, and W. R. Jarvis. 1995. Postoperative infections traced to contamination of an intravenous anesthetic, propofol. *N. Engl. J. Med.* 333:147–154.
- Berry, N., I. Hassan, S. Majumdar, A. Vardhan, A. McEwen, and R. Gokal. 2004. *Bacillus circulans* peritonitis in a patient treated with CAPD. *Perit. Dial. Int.* 24:488–489.
- Bishop, A. L., and R. M. Davis. 1990. Internal decay of onions caused by *Enterobacter cloacae*. *Plant Dis.* 74:692–694.
- Blanchard, R. O., and T. A. Tattar. 1981. *Field and Laboratory Guide to Tree Pathology*. Academic Press, New York, N.Y.
- Blodgett, J. T., and W. J. Swart. 2002. Infection, colonization and disease of *Amaranthus hybridus* leaves by the *Alternaria tenuissima* group. *Plant Dis.* 86:1199–1205.
- Boby, V., U., and D. J. Bagyaraj. 2003. Biological control of root-rot of *Coleus forskohlii* Briq. using microbial inoculants. *World J. Microbiol.* 19:175–180.
- Bottalico, A., and G. Perrone. 2002. Toxigenic *Fusarium* species and mycotoxins associated with head blight in small-grain cereals in Europe. *Eur. J. Plant Pathol.* 108:611–624.
- Boyd, A. S., B. Wiser, H. Sams, and L. King. 2003. Gangrenous cutaneous mucormycosis in a child with a solid organ transplant: a case report and review of the literature. *Pediatr. Dermatol.* 20:411–415.
- Brazier, J. S., M. Gal, V. Hall, and T. E. Morris. 2004. Outbreak of *Clostridium histolyticum* infections in injecting drug users in England and Scotland. *Eurosurveillance Mon.* 9:15–16. [Online.] <http://www.eurosurveillance.org/em/v09n09/0909-221.asp>.
- Brenner, D. J., J. R. Rodrigues Neto, A. G. Steigerwalt, and C. F. Robbs. 1994. "*Erwinia nalandii*" is a subjective synonym of *Erwinia persicinus*. *Int. J. Syst. Bacteriol.* 44:282–284.
- Bridges, C. H. 1957. Maduromycotic mycetomas in animals: *Curvularia geniculata* as an etiologic agent. *Am. J. Pathol.* 33:411–427.
- Brown, R. L., T. E. Cleveland, P. J. Cotty, and J. E. Mellon. 1992. Spread of *Aspergillus flavus* in cotton bolls, decay of intercarpellary membranes and production of fungal pectinases. *Phytopathology* 82:462–467.
- Bruton, B. D., F. Mitchell, J. Fletcher, S. D. Pair, A. Wayadande, U. Melcher, J. Brady, B. Bextine, and T. W. Popham. 2003. *Serratia marcescens*, a phloem-colonizing, squash bug-transmitted bacterium: causal agent of cucurbit yellow vine disease. *Plant Dis.* 87:937–944.
- Bugos, R. C., J. B. Sutherland, and J. H. Adler. 1988. Phenolic compound utilization by the soft rot fungus *Lecythophora hoffmannii*. *Appl. Environ. Microbiol.* 54:1882–1885.
- Burgess, L. W., and C. M. Liddell. 1983. *Laboratory Manual for Fusarium Research*. University of Sydney Press, Sydney, Australia.
- Burkholder, W. H. 1950. Sour skin, a bacterial rot of onion bulbs. *Phytopathology* 40:115–117.
- Bushelman, S. J., J. P. Callen, D. N. Roth, and L. M. Cohen. 1995. Disseminated *Fusarium solani* infection. *J. Am. Acad. Dermatol.* 32:346–351.
- Byrd, R. P., T. M. Roy, C. L. Fields, and J. A. Lynch. 1992. *Paecilomyces variotii* pneumonia in a patient with diabetes mellitus. *J. Diabet. Complicat.* 6:150–153.
- Camin, A. M., C. Michelet, T. Langanay, C. de Place, S. Chevrier, E. Gueho, and C. Guiguen. 1999. Endocarditis due to *Fusarium dimerum* four years after coronary artery bypass grafting. *Clin. Infect. Dis.* 28:150.

- Cao, H., R. L. Baldini, and L. G. Rahme. 2001. Common mechanisms for pathogens of plants and animals. *Annu. Rev. Phytopathol.* 39:259–284.
- Carey, J., M. Motyl, and D. C. Perlman. 2001. Catheter-related bacteremia due to *Streptomyces* in a patient receiving holistic infusions. *Emerg. Infect. Dis.* 7:1043–1045.
- Carter, E., and C. Boudreaux. 2004. Fatal cerebral phaeohyphomycosis due to *Curvularia lunata* in an immunocompetent patient. *J. Clin. Microbiol.* 42:5419–5423.
- Centers for Disease Control and Prevention and National Institutes of Health. 1999. *Biosafety in Microbiological and Biomedical Laboratories*, 4th ed. U.S. Government Printing Office, Washington, D.C.
- Chalet, M., D. H. Howard, M. R. McGinnis, and I. Zapatero. 1986. Isolation of *Bipolaris australiensis* from a lesion of viral vesicular dermatitis on the scalp. *J. Med. Vet. Mycol.* 24:461–465.
- Chandra, S., and A. Woodgyer. 2002. Primary cutaneous zygomycosis due to *Mucor circinelloides*. *Australas. J. Dermatol.* 43:39–42.
- Chase, A. R., J. W. Miller, and J. B. Jones. 1984. Leaf spot and blight of *Asplenium nidus* caused by *Pseudomonas gladioli*. *Plant Dis.* 68:344–347.
- Chuenchitt, S., W. Dhirabhava, S. Karnjanarat, D. Buangsuwon, and T. Uematsu. 1983. A new bacterial disease on orchids *Dendrobium* sp. caused by *Pseudomonas gladioli*. *Kasetsart J. Nat. Sci.* 17:26–36.
- Cother, E. J., B. Darbyshire, and J. Brewer. 1976. *Pseudomonas aeruginosa*: cause of internal brown rot of onion. *Phytopathology* 66:828–834.
- Courtney, J. M., K. E. Dunbar, A. McDowell, J. E. Moore, T. J. Warke, M. Stevenson, and J. S. M. Elborn. 2004. Clinical outcome of *Burkholderia cepacia* complex infection in cystic fibrosis adults. *J. Cyst. Fibros.* 3:93–98.
- Dantas, S., S. Oliveira, and S. Michereff. 2003. Post harvest fungal diseases in papaya and orange marketed in the distribution centre of Recife. *Fitopatol. Bras.* 28:528–533.
- De Boeck, K., A. Malfroot, L. Van Schil, P. Lebecque, C. Knoop, J. R. Govan, C. Doherty, S. Laevens, and P. Vandamme, on behalf of Belgian *Burkholderia cepacia* Study Group. 2004. Epidemiology of *Burkholderia cepacia* complex colonization in cystic fibrosis patients. *Eur. Respir. J.* 23:851–856.
- Denton, M., and K. G. Kerr. 1998. Microbiological and clinical aspects of infection associated with *Stenotrophomonas maltophilia*. *Clin. Microbiol. Rev.* 11:57–80.
- Deva, A. K., and K. G. Narayan. 1989. Enterotoxigenicity of *Bacillus circulans*, *B. coagulans* and *B. stearothermophilus*. *Indian J. Comp. Microbiol. Immunol. Infect. Dis.* 10:80–87.
- Dhingra, O. D., and J. B. Sinclair. 1995. *Basic Plant Pathology Methods*, 2nd ed. CRC Press, Inc., Boca Raton, Fla.
- Dickman, M. D., and A. M. Alvarez. 1983. Latent infection of papaya caused by *Colletotrichum gloeosporioides*. *Plant Dis.* 67:748–750.
- Dignani, M. C., and E. Anaissie. 2004. Human fusariosis. *Clin. Microbiol. Infect.* 10(Suppl. 1):67–75.
- Dillard, H. R., and A. C. Cobb. 1997. Disease progress of black dot on tomato roots and reduction in incidence with foliar applied fungicides. *Plant Dis.* 81:1439–1442.
- Dimitrov, D. S. 2004. Virus entry: molecular mechanisms and biomedical applications. *Nat. Rev. Microbiol.* 2: 109–122.
- Dixon, D. M., M. M. McNeil, M. L. Cohen, B. G. Gellin, and J. R. La Montagne. 1996. Fungal infections: a growing threat. *Public Health Rep.* 111:226–235.
- Dong, Y., M. K. Chelius, S. Brisse, N. Kozyrovska, R. Podschun, and E. W. Triplett. 2003. Comparisons between two *Klebsiella*: the plant endophyte *K. pneumoniae* 342 and a clinical isolate, *K. pneumoniae* MGH78578. *Symbiosis* 35:247–259.
- Duffill, M. B., and K. E. Coley. 1993. Cutaneous phaeohyphomycosis due to *Alternaria alternata* responding to itraconazole. *Clin. Exp. Dermatol.* 18:156–158.
- Dunne, W. M., Jr., J. Tillman, and J. C. Murray. 1993. Recovery of a strain of *Agrobacterium radiobacter* with a mucoid phenotype from an immunocompromised child with bacteremia. *J. Clin. Microbiol.* 31:2541–2543.
- Ebright, J. R., P. H. Chandrasekar, S. Marks, M. R. Fairfax, A. Aneziokoro, and M. R. McGinnis. 1999. Invasive sinusitis and cerebritis due to *Curvularia clavata* in an immunocompetent adult. *Clin. Infect. Dis.* 28:687–689.
- Edgar, J. 2004. Future impact of food safety issues on animal production and trade: implications for research. *Austr. J. Exp. Agr.* 44:1073–1076.
- Edmond, M. B., S. A. Riddler, C. M. Baxter, B. M. Wicklund, and A. W. Pasculle. 1993. *Agrobacterium radiobacter*: a recently recognized opportunistic pathogen. *Clin. Infect. Dis.* 16:388–391.
- Eisen, D. P., and J. Robson. 2004. Complete resolution of pulmonary *Rhizopus oryzae* infection with itraconazole treatment: more evidence of the utility of azoles for zygomycosis. *Mycoses* 47:159–162.
- Engelhard, D. 1998. Bacterial and fungal infections in children undergoing bone marrow transplantation. *Bone Marrow Transplant.* 21(Suppl. 2):S78–S80.
- Fahy, P. C., and G. J. Persley. 1983. *Plant Bacterial Diseases. A Diagnostic Guide*. Academic Press, New York, N.Y.
- Fata, F., S. Chittivelu, S. Tessler, and Y. Kupfer. 1996. Gas gangrene of the arm due to *Enterobacter cloacae* in a neutropenic patient. *South. Med. J.* 89:1095–1096.
- Fischl, G., L. Szunics, and J. Bakonyi. 1993. Black points in wheat grains. *Noenmytermeles* 42:419–429.
- Flanagan, K. L., and A. D. Bryceson. 1997. Disseminated infection due to *Bipolaris australiensis* in a young immunocompetent man: case report and review. *Clin. Infect. Dis.* 25:311–313.
- Frederickson, D. E., E. S. Monyo, S. B. King, and G. N. Odvody. 1997. A disease of pearl millet in Zimbabwe caused by *Pantoea agglomerans*. *Plant Dis.* 81:959.
- Funke, G., G. Haase, N. Schnitzler, N. Schrage, and R. R. Reinert. 1997. Endophthalmitis due to *Microbacterium* species: case report and review of *Microbacterium* infections. *Clin. Infect. Dis.* 24:713–716.
- Geiser, D. M., J. W. Dörner, B. W. Horn, and J. W. Taylor. 2000. The phylogenetics of mycotoxin and sclerotium production in *Aspergillus flavus* and *Aspergillus oryzae*. *Fungal Genet. Biol.* 31:169–179.
- Georg, L. K. 1964. *Curvularia geniculata*, a cause of mycotic keratitis. *J. Med. Assoc. State Ala.* 33:234–236.
- Geraldi, M. A. P., M. F. Ito, A. Ricci, Jr., F. O. Paradela, and H. Nagal. 1980. *Chaetomium globosum* Kunze, causal agent of a new tomato disease. *Summa Phytopathol.* 6:79–84.
- Ghosh, K., S. Daar, D. Hiwase, and N. Nursat. 2000. Primary *Pseudomonas* meningitis in an adult, splenectomized, multitransfused thalassaemia major patient. *Haematologia* 30:69–72.
- Gibbs, M. J., and G. Weiller. 1999. Evidence that a plant virus switched hosts to infect a vertebrate and then recombined with a vertebrate-infecting virus. *Proc. Natl. Acad. Sci. USA* 96:8022–8027.

- Gill, W. M., and A. L. J. Cole. 1992. Cavity disease of *Agaricus bitorquis* caused by *Pseudomonas cepacia*. *Can. J. Microbiol.* 38:394-397.
- Glick, B. R. 2004. Teamwork in phytoremediation. *Nat. Biotechnol.* 22:526-527.
- Godoy, P., F. Nunes, V. Silva, J. Tomimori-Yamashita, L. Zaror, and O. Fischman. 2004. Onychomycosis caused by *Fusarium solani* and *Fusarium oxysporum* in São Paulo, Brazil. *Mycopathologia* 157:287-290.
- Gonzalez, A., A. del Palacio, M. S. Cuetara, C. Gomez, E. Carabias, and Q. Malo. 1996. Zygomycosis: review of 16 cases. *Enferm. Infecc. Microbiol. Clin.* 14:233-239.
- Gorbalenya, A. E., A. P. Donchenko, V. M. Blinov, and E. V. Koonin. 1989. Cysteine proteases of positive strand RNA viruses and chymotrypsin-like serine proteases. A distinct protein superfamily with a common structural fold. *FEBS Lett.* 243:103-114.
- Goss, C. H., N. Mayer-Hamblett, M. L. Aitken, G. D. Rubenfeld, and B. W. Ramsey. 2004. Association between *Stenotrophomonas maltophilia* and lung function in cystic fibrosis. *Thorax* 59:955-959.
- Gour, H. N., and H. C. Dube. 1975. Production of pectic enzymes by *Curvularia lunata* causing leaf spot of cotton. *Proc. Indian. Natl. Sci. Acad. Part B.* 41:480-485.
- Govan, J. R., J. E. Hughes, and P. Vandamme. 1996. *Burkholderia cepacia*: medical, taxonomic and ecological issues. *J. Med. Microbiol.* 45:395-407.
- Govan, J. R., and P. Vandamme. 1998. Agricultural and medical microbiology: a time for bridging gaps. *Microbiology* 144:2373-2375.
- Graves, M., T. Robin, A. M. Chipman, J. Wong, S. Khashe, and J. M. Janda. 1997. Four additional cases of *Burkholderia gladioli* infection with microbiological correlates and review. *Clin. Infect. Dis.* 25:838-842.
- Grimont, P. A. D., F. Grimont, and M. P. Starr. 1979. *Serratia ficaria* sp. nov., a bacterial species associated with Smyrna figs and the fig wasp *Blastophaga psenes*. *Curr. Microbiol.* 2:277-282.
- Guarro, J., T. Akiti, R. Almada-Horta, L. A. Leite-Filho, J. Gene, S. Ferreira-Gomes, C. Aguilar, and M. Ortoneda. 1999. Mycotic keratitis due to *Curvularia senegalensis* and *in vitro* antifungal susceptibilities of *Curvularia* spp. *J. Clin. Microbiol.* 37:4170-4173.
- Gugnani, H. C., C. N. Okeke, and A. Sivanesan. 1990. *Curvularia clavata* as an etiologic agent of human skin infection. *Lett. Appl. Microbiol.* 10:47-49.
- Gupta, V., R. Chawla, and S. Sen. 2001. *Aureobasidium pullulans* scleritis following keratoplasty: a case report. *Ophthalmic Surg. Lasers* 32:481-482.
- Gvozdiak, R. I., S. F. Khodos, and V. V. Lipshits. 1976. Biological properties of *Clostridium butyricum* V. phytopathogenicum var. nova, the causative agent of horn-beam disease. *Mikrobiol. Zh.* 38:288-292. (In Russian.)
- Gyaneshwar, P., E. K. James, N. Mathan, P. M. Reddy, B. Reinhold-Hurek, and J. K. Ladha. 2001. Endophytic colonization of rice by a diazotrophic strain of *Serratia marcescens*. *J. Bacteriol.* 183:2634-2645.
- Hall, N., J. P. R. Keon, and J. A. Hargreaves. 1999. A homologue of a gene implicated in the virulence of human fungal diseases is present in a plant fungal pathogen and is expressed during infection. *Physiol. Mol. Plant Pathol.* 55:69-73.
- Hammouda, A. M. 1992. A new leaf spot of pepper caused by *Cladosporium oxysporum*. *Plant Dis.* 76:536-537.
- Hampton, R., E. Ball, and S. DeBoer. 1990. Serological Methods for Detection of Viral and Bacterial Plant Pathogens: a Laboratory Manual. APS Press, Inc., St. Paul, Minn.
- Hao, M. V., D. J. Brenner, A. G. Steigerwalt, Y. Kosako, and K. Komugata. 1990. *Erwinia persicinus*, a new species isolated from plants. *Int. J. Syst. Bacteriol.* 40:379-383.
- Hattori, N., M. Adachi, T. Kaneko, M. Shimosuma, M. Ichinohe, and K. Iozumi. 2000. Case report: onychomycosis due to *Chaetomium globosum* successfully treated with itraconazole. *Mycoses* 43:89-92.
- Heimann, M. F., and D. M. Boone. 1983. Raspberry (*Rubus*) disorders: cane blight and spur blight [*Leptosphaeria coniothyrium*, *Coniothyrium fuckelii*, Wisconsin]. Publ. Coop. Ext. Programs University of Wisconsin. Ext.: The Programs.
- Hickey, K. D. 1986. *Methods for Evaluating Pesticides for Control of Plant Pathogens*. APS Press, Inc., St. Paul, Minn.
- Hinton, D. M., and C. W. Bacon. 1995. *Enterobacter cloacae* is an endophytic symbiont of corn. *Mycopathologia* 129:117-125.
- Hohn, T., and J. Fütterer. 1997. The proteins and function of plant pararetroviruses: knowns and unknowns. *Crit. Rev. Plant Sci.* 16:133-161.
- Holmes, A., R. Nolan, R. Taylor, R. Finley, M. Riley, R. Z. Jiang, S. Steinbach, and R. Goldstein. 1999. An epidemic of *Burkholderia cepacia* transmitted between patients with and without cystic fibrosis. *J. Infect. Dis.* 179:1197-1205.
- Honda, Y., M. Z. Rahman, S. Z. Islam, and N. Muroguchi. 2001. Leaf spot disease of broad bean caused by *Alternaria tenuissima* in Japan. *Plant Dis.* 85:95.
- Horner, W. E., A. Helbling, J. E. Salvaggio, and S. B. Lehrer. 1995. Fungal allergens. *Clin. Microbiol. Rev.* 8:161-179.
- Howard, C. M., and E. E. Albregts. 1973. A strawberry fruit rot caused by *Alternaria tenuissima*. *Phytopathology* 63:938-939.
- Howard, F. M., J. M. Bradley, D. M. Flynn, P. Noone, and M. Szawatkowski. 1977. Outbreak of necrotising enterocolitis caused by *Clostridium butyricum*. *Lancet* 310:1099-1102.
- Hsueh, P. R., L. J. Teng, H. J. Pan, Y. C. Chen, C. C. Sun, S. W. Ho, and K. T. Luh. 1998. Outbreak of *Pseudomonas fluorescens* bacteremia among oncology patients. *J. Clin. Microbiol.* 36:2914-2917.
- Hueck, C. J. 1998. Type III protein secretion systems in bacterial pathogens of animals and plants. *Microbiol. Mol. Biol. Rev.* 62:379-433.
- Hulse, M., S. Johnson, and P. Ferrieri. 1993. *Agrobacterium* infections in humans: experience at one hospital and review. *Clin. Infect. Dis.* 16:112-117.
- Ito, M. F., F. Paradela, J. Soave, and M. H. Sugimori. 1979. Leaf spot caused in maize (*Zea mays* L.) by *Curvularia lunata*. *Summa Phytopathol.* 5:181-184.
- James, D., and C. Upton. 2002. Nucleotide sequence analysis of RNA-2 of a flat apple isolate of Cherry rasp leaf virus with regions showing greater identity to animal picornaviruses than to related plant viruses. *Arch. Virol.* 147:1631-1641.
- Jochimsen, E. M., C. Frenette, M. Delorme, M. Arduino, S. Aguero, L. Carson, J. Ismail, S. Lapierre, E. Czyziw, J. I. Tokars, and W. R. Jarvis. 1998. A cluster of bloodstream infections and pyrogenic reactions among hemodialysis patients traced to dialysis machine waste-handling option units. *Am. J. Nephrol.* 18:485-489.
- Johansen, H. K., T. A. Kovesi, C. Koch, M. Corey, N. Hoiby, and H. Levinson. 1998. *Pseudomonas aeruginosa* and *Burkholderia cepacia* infection in cystic fibrosis

- patients treated in Toronto and Copenhagen. *Pediatr. Pulmonol.* 26:89-96.
- Johnson, K. B., and V. O. Stockwell. 1998. Management of fire blight: a case study in microbial ecology. *Annu. Rev. Phytopathol.* 36:227-248.
- Jones, D. R., and E. O. Lomeiro. 2000. Pseudostem heart rot, p. 166-167. In D. R. Jones (ed.), *Diseases of Banana, Abaca and Enset*. CABI Publishing, Wallingford, United Kingdom.
- Jones, D. R., and R. H. Stover. 2000. Fungal root rot, p. 162-163. In D. R. Jones (ed.), *Diseases of Banana, Abaca and Enset*. CABI Publishing, Wallingford, United Kingdom.
- Kaczmarek, E. B., J. A. Liu Yin, J. A. Tooth, E. M. Love, and I. W. Delamore. 1986. Systemic infection with *Aureobasidium pullulans* in a leukaemic patient. *J. Infect.* 13:289-291.
- Kahn, R. P., and S. B. Mathur (ed.). 1999. *Containment Facilities and Safeguards for Exotic Plant Pathogens and Pests*. APS Press, St. Paul, Minn.
- Kaku, H. 2004. Histopathology of red stripe of rice. *Plant Dis.* 88:1304-1309.
- Kang, C. I., S. H. Kim, D. M. Kim, W. B. Park, K. D. Lee, H. B. Kim, M. D. Oh, E. C. Kim, and K. W. Choe. 2004. Risk factors for and clinical outcomes of bloodstream infections caused by extended-spectrum beta-lactamase-producing *Klebsiella pneumoniae*. *Infect. Control Hosp. Epidemiol.* 25:860-867.
- Kantarcioğlu, A. S., G. Hatemi, A. Yucel, G. S. De Hong, and N. M. Mandel. 2003. *Paecilomyces variotii* central nervous system infection in a patient with cancer. *Mycoses* 46:45-50.
- Karasev, A. V., S. S. Han, and T. Iwanami. 2001. Satsuma dwarf and related viruses belong to a new lineage of plant picorna-like viruses. *Virus Genes* 23:45-52.
- Khoa, L. V., K. Hatai, and T. Aoki. 2004. *Fusarium incarnatum* isolated from black tiger shrimp, *Penaeus monodon* Fabricius, with black gill disease cultured in Vietnam. *J. Fish Dis.* 27:507-515.
- Kiehn, T. E., B. Polsky, E. Punithalingam, F. F. Edwards, A. E. Brown, and D. Armstrong. 1987. Liver infection caused by *Coniothyrium fuckelii* in a patient with acute myelogenous leukemia. *J. Clin. Microbiol.* 25:2410-2412.
- Kishun, R., and J. Swarup. 1981. Growth studies on *Pseudomonas gladioli* pv. *allicola* pathogenic to onion. *Indian J. Mycol. Plant Pathol.* 11:247-250.
- Klement, Z., K. Rudolph, and D. C. Sands (ed.). 1990. *Methods in Phytobacteriology*. Akademiai Kiado, Budapest, Hungary.
- Kliejunas, J. T., J. R. Allison, A. H. McCain, and R. S. Smith, Jr. 1985. *Phoma* blight of fir and Douglas fir seedlings in a California nursery. *Plant Dis.* 69:773-775.
- Ko, W. H., I. T. Wang, and P. J. Ann. 2004. *Lasiodiplodia theobromae* as a causal agent of kumquat dieback in Taiwan. *Plant Dis.* 88:1383.
- Kore, S. S., and V. P. Bhide. 1976. A first report of *Curvularia brachyspora* Boedijn inciting leaf-spot disease of rose. *Curr. Sci.* 45:74.
- Kratz, A., D. Greenberg, Y. Barki, E. Cohen, and M. Lifshitz. 2003. *Pantoea agglomerans* as a cause of septic arthritis after palm tree thorn injury; case report and literature review. *Arch. Dis. Child.* 88:542-544.
- Krysinska-Traczyk, E., and J. Dutkiewicz. 2000. *Aspergillus candidatus*: a respiratory hazard associated with grain dust. *Ann. Agric. Environ. Med.* 7:101-109.
- Lakshmanan, P. 1992. Sheath rot of rice incited by *Curvularia lunata* in Tamal Nadu, India. *Trop. Pest Manage.* 38:107.
- Lal, S., and H. S. Tripathi. 1977. Host range of *Curvularia pallescens*, the incitant of leaf spot of maize. *Indian J. Mycol. Plant Pathol.* 7:92-93.
- Latham, R. H. 2000. *Bipolaris spicifera* meningitis complicating a neurosurgical [sic] procedure. *Scand. J. Infect. Dis.* 32:102-103.
- Lau, S. M., W. L. Yu, and J. H. Wang. 1999. Cardiac cirrhosis with cellulitis caused by *Burkholderia cepacia* bacteremia. *Clin. Infect. Dis.* 29:447-448.
- Laverde, S., L. H. Moncada, A. Restrepo, and C. L. Vera. 1973. Mycotic keratitis—5 cases caused by unusual fungi. *Sabouraudia* 11:119-123.
- Leach, C. M., and M. Tulloch. 1972. World-wide occurrence of the suspected mycotoxin producing fungus *Drechslera biseptata* with grass seed. *Mycologia* 64:1357-1359.
- Leary, J. V., and W. W. W. Chun. 1989. Pathogenicity of *Bacillus circulans* to seedlings of date palm (*Phoenix dactylifera*). *Plant Dis.* 73:353-354.
- Lelliot, R. A., and D. E. Stead. 1987. *Methods for the Diagnosis of Bacterial Diseases of Plants*. Blackwell Scientific Publications, Boston, Mass.
- Li, Z. X., Z. S. Bian, H. P. Zheng, Y. S. Yue, J. Y. Yao, Y. P. Gong, M. Y. Cai, and X. Z. Dong. 1990. First isolation of *Xanthomonas campestris* from the blood of a Chinese woman. *Chin. Med. J.* 103:435-439.
- Linares, G., P. A. McGarry, and R. D. Baker. 1971. Solid solitary aspergillotic granuloma of the brain. Report of a case due to *Aspergillus candidus* and review of the literature. *Neurology* 21:177-184.
- Lombardi, G., F. Luzzaro, J. Docquier, M. L. Riccio, M. Perilli, A. Coli, G. Amicosante, G. M. Rossolini, and A. Toniolo. 2002. Nosocomial infections caused by multidrug-resistant isolates of *Pseudomonas putida* producing VIM-1 metallo- β -lactamase. *J. Clin. Microbiol.* 40:4051-4055.
- Lucas, J. A., and C. H. Dickinson. 1998. *Plant Pathology and Plant Pathogens*, 3rd ed. Blackwell Science, Oxford, England.
- Lukezic, F. L., D. C. Hildebrand, M. N. Schroth, and P. A. Shinde. 1982. Association of *Serratia marcescens* with crown rot of alfalfa in Pennsylvania. *Phytopathology* 72:714-718.
- Mandokhot, A. M., and K. C. B. Chaudhary. 1972. A new leaf spot of maize incited by *Curvularia clavata*. *Neth. J. Plant Pathol.* 78:65-68.
- Mangiaterra, M., G. Giusiano, G. Smilasky, L. Zamar, G. Amado, and C. Vincentin. 2001. Keratomycosis caused by *Cylindrocarpum lichenicola*. *Med. Mycol.* 39:143-145.
- Marcus, L., H. F. Vismer, H. J. van der Hoven, E. Gove, and P. Meewes. 1992. Mycotic keratitis caused by *Curvularia brachyspora* (Boedijn) [sic]. A report of the first case. *Mycopathologia* 119:29-33.
- Marriott, D. J., K. H. Wong, E. Aznar, J. L. Harkness, D. A. Cooper, and D. Muir. 1997. *Scytalidium dimidiatum* and *Lecythophora hoffmannii*: unusual causes of fungal infections in a patient with AIDS. *J. Clin. Microbiol.* 35:2949-2952.
- Maslen, M. M., T. Collis, and R. Stuart. 1996. *Lasiodiplodia theobromae* isolated from a subcutaneous abscess in a Cambodian immigrant to Australia. *Med. Mycol.* 34:279-283.
- Matteson Heidenreich, M. C., M. R. Corral-Garcia, E. A. Momol, and T. J. Burr. 1997. Russet of apple fruit caused by *Aureobasidium pullulans* and *Rhodotorula glutinis*. *Plant Dis.* 81:337-342.
- McGinnis, M. R., G. Campbell, W. K. Gourley, and H. L. Lucia. 1992. Phaeohyphomycosis caused by *Bipolaris*

- spicifera*: an informative case. *Eur. J. Epidemiol.* 8:383–386.
- McKeen, W. E. 1989. *Blue Mold of Tobacco*. American Phytopathology Society Press, St. Paul, Minn.
- Melgosa Hijosa, M., M. C. Ramos Lopez, P. Ruiz Almagro, A. Fernandez Escribano, and A. Luque de Pablos. 1997. *Agrobacterium radiobacter* peritonitis in a Down's syndrome child maintained on peritoneal dialysis. *Perit. Dial. Int.* 17:515.
- Meredith, D. S. 1963. Some graminicolous fungi associated with spotting of banana leaves in Jamaica. *Ann. App. Biol.* 51:371–378.
- Mettenleiter, T. C. 2002. Brief overview on cellular virus receptors. *Virus Res.* 82:3–8.
- Michailides, T. J., D. P. Morgan, and K. V. Subbarao. 1996. Fig endosepsis: an old disease still a dilemma for California growers. *Plant Dis.* 80:828–841.
- Milholland, R. D. 1995. *Alternaria* leaf spot and fruit rot, p. 18. In F. L. Caruso and D. C. Ramsdell (ed.), *Compendium of Blueberry and Cranberry Diseases*. APS Press, St. Paul, Minn.
- Miller, J. M., C. Novy, and M. Hiott. 1996. Case of bacterial endophthalmitis caused by an *Agrobacterium radiobacter*-like organism. *J. Clin. Microbiol.* 34:3212–3213.
- Mishra, G. S., N. Mehta, and M. Pal. 2004. Chronic bilateral otomycosis caused by *Aspergillus niger*. *Mycoses* 47:82–84.
- Moore, L. W., and G. Warren. 1979. *Agrobacterium radiobacter* strain 84 and biological control of crown gall. *Annu. Rev. Phytopathol.* 17:163–179.
- Morton, S. J., K. Midthun, and W. G. Merz. 1986. Granulomatous encephalitis caused by *Bipolaris hawaiiensis*. *Arch. Pathol. Lab. Med.* 110:1183–1185.
- Muchovej, J. J., and H. B. Couch. 1987. Colonization of bentgrass turf by *Curvularia lunata* after leaf clipping and heat stress. *Plant Dis.* 71:873–875.
- Mullen, J. M., C. H. Gilliam, A. K. Hagan, and G. Morgan Jones. 1991. Canker of dogwood caused by *Lasiodiplodia theobromae*, a disease influenced by drought stress or cultivar selection. *Plant Dis.* 75:886–889.
- Murdoch, C. W., and R. J. Campana. 1983. Bacterial species associated with wetwood of elm. *Phytopathology* 73:1270–1273.
- Murray, C. K., M. L. Beckius, and K. McAllister. 2003. *Fusarium proliferatum* superficial suppurative thrombophlebitis. *Mil. Med.* 168:426–427.
- Mycology Online. University of Adelaide, Adelaide, Australia. [http://www.mycology.adelaide.edu.au/Fungal_Descriptions/Hyphomycetes_\(dematiaceous\)/Drechslera/index.html](http://www.mycology.adelaide.edu.au/Fungal_Descriptions/Hyphomycetes_(dematiaceous)/Drechslera/index.html)
- Naiker, S., and B. Odhav. 2004. Mycotic keratitis: profile of *Fusarium* species and their mycotoxins. *Mycoses* 47:50–56.
- Namdari, H., S. Hamzavi, and R. R. Peairs. 2003. *Rhizobium* (*Agrobacterium*) *radiobacter* identified as a cause of chronic endophthalmitis subsequent to cataract extraction. *J. Clin. Microbiol.* 41:3998–4000.
- National Institutes of Health. 2002. *NIH Guidelines for Research Involving Recombinant DNA Molecules*. NIH Guidelines 59 CFR 34496 (July 5, 1994), as amended. [Online. The current amended version of the NIH guidelines can be accessed at <http://www4.od.nih.gov/oba/rac/guidelines/guidelines.html>.]
- Nault, L. R. 1997. Arthropod transmission of plant viruses: a new synthesis. *Ann. Entomol. Soc. Am.* 90:521–542.
- Nelson, K. E., C. Weinl, T. Paulsen, R. J. Dodson, H. Hilbert, V. A. P. Martins dos Santos, D. E. Fouts, S. R. Gill, M. Pop, M. Holmes, L. Brinkac, M. Beanan, R. T. DeBoy, S. Daugherty, J. Kolonay, R. Madupu, W. Nelson, O. White, J. Peterson, H. Khouri, I. Hance, P. C. Lee, E. Holtzapple, D. Scanlan, K. Tran, A. Moazzez, T. Utterback, M. Rizzo, K. Lee, D. Kosack, D. Moestl, H. Wedler, J. Lauber, D. Stjepandic, J. Hoheisel, M. Straetz, S. Heim, C. Kiewitz, J. Eisen, K. N. Timmis, A. Dusterhöft, B. Tümmeler, and C. M. Fraser. 2002. Complete genome sequence and comparative analysis of the metabolically versatile *Pseudomonas putida*. *Environ. Microbiol.* 4:799–808.
- Nishijima, K. A., A. M. Alvarez, P. R. Hepperly, M. H. Shintaku, L. M. Keith, D. M. Sato, B. C. Bushe, J. W. Armstrong, and F. T. Zee. 2004. Association of *Enterobacter cloacae* with rhizome rot of edible ginger in Hawaii. *Plant Dis.* 88:1318–1327.
- Nyvall, R. F. 1979. *Field Crop Diseases Handbook*. AVI Publishing Company, Inc., Westport, Conn.
- Ogden, P. E., D. L. Hurley, and P. T. Cain. 1992. Fatal fungal endarteritis caused by *Bipolaris spicifera* following replacement of the aortic valve. *Clin. Infect. Dis.* 14:596–598.
- O'Hara, C. M., A. G. Steigerwalt, B. C. Hill, J. M. Miller, and D. J. Brenner. 1998. First report of a human isolate of *Erwinia persicinus*. *J. Clin. Microbiol.* 36:248–250.
- Ono, M., C. Nishigori, C. Tanaka, S. Tanaka, M. Tsuda, and Y. Miyachi. 2004. Cutaneous alternariosis in an immunocompetent patient: analysis of the internal transcribed spacer region of rDNA and Brm2 of isolated *Alternaria alternata*. *Br. J. Dermatol.* 150:773–775.
- O'Quinn, R. P., J. L. Hoffmann, and A. S. Boyd. 2001. *Colletotrichum* species as emerging opportunistic fungal pathogens: a report of 3 cases of phaeohyphomycosis and review. *J. Am. Acad. Dermatol.* 45:56–61.
- O'Shaughnessy, E. M., G. N. Forrest, and T. J. Walsh. 2003. Invasive aspergillosis in patients with hematologic malignancies: recent advances and new challenges. *Curr. Treat. Opt. Infect. Dis.* 5:507–515.
- Ostrowsky, B. E., C. Whitener, H. K. Bredenberg, L. A. Carson, S. Holt, L. Hutwagner, M. J. Arduino, and W. R. Jarvis. 2002. *Serratia marcescens* bacteremia traced to an infused narcotic. *N. Engl. J. Med.* 346:1529–1537.
- Pathengay, A., B. Raju, S. Sharma, T. Das, and Endophthalmitis Research Group. 2005. Recurrent endophthalmitis caused by *Burkholderia cepacia*. *Eye* 19:358–359. (First published 30 July 2004; doi:10.1038/sj.eye.6701501.)
- Phipps, P. M., and D. M. Porter. 1998. Collar rot of peanut caused by *Lasiodiplodia theobromae*. *Plant Dis.* 82:1205–1209.
- Pitt, J. I., S. K. Dyer, and S. McCammon. 1991. Systemic invasion of developing peanut plants by *Aspergillus flavus*. *Lett. Appl. Microbiol.* 13:16–20.
- Plotkin, G. R. 1980. *Agrobacterium radiobacter* prosthetic valve endocarditis. *Ann. Intern. Med.* 93:839–840.
- Pontón, J., R. Ruchel, K. V. Clemons, D. C. Colemans, R. Grillot, J. Guarro, D. Aldebert, P. Ambroise-Thomas, J. Cano, A. J. Carrillo-Munoz, J. Gene, C. Pinel, D. A. Stevens, and D. J. Sullivan. 2000. Emerging pathogens. *Med. Mycol.* 38(Suppl. 1):225–236.
- Pratt, R. G. 2001. Occurrence and virulence of *Bipolaris hawaiiensis* on bermudagrass (*Cynodon dactylon*) on poultry waste application sites in Mississippi. *Plant Dis.* 85:1206.
- Prince, S. E., K. A. Dominger, B. A. Cunha, and N. C. Klein. 1997. *Klebsiella pneumoniae* pneumonia. *Heart Lung* 26:413–417.
- Procop, G. W., and G. D. Roberts. 1998. Laboratory methods in basic mycology, p. 871–961. In B. A. Forbes,

- D. F. Sahm, and A. S. Weissfeld (ed.), *Bailey and Scott's Diagnostic Microbiology*, 10th ed. Mosby, St. Louis, Mo.
- Punja, Z. K. 1997. Comparative efficacy of bacteria, fungi and yeasts as biological control agents for diseases of vegetable crops. *Can. J. Plant Pathol.* 19:315-323.
- Raabe, R. D., L. L. Conners, and A. P. Martinez. 1981. Checklist of plant disease in Hawaii: including records of microorganisms, principally fungi, found in the state. Information text series 022. Hawaii Institute of Tropical Agriculture and Human Resources, University of Hawaii at Manoa.
- Rahme, L. G., E. J. Stevens, S. F. Wolfort, J. Shao, R. G. Tompkins, and F. M. Ausubel. 1995. Common virulence factors for bacterial pathogenicity in plants and animals. *Science* 268:1899-1902.
- Ramirez, F. C., Z. A. Saeed, R. O. Darouiche, R. M. Shawar, and B. Yoffe. 1992. *Agrobacterium tumefaciens* peritonitis mimicking tuberculosis. *Clin. Infect. Dis.* 15:938-940.
- Rao, G. P., S. P. Singh, and M. Singh. 1992. Two alternative hosts of *Curvularia pallescens*, the leaf spot causing fungus of sugarcane. *Trop. Pest Manag.* 38:218.
- Rao, K., and V. Saha. 2000. Medical management of *Aspergillus flavus* endocarditis. *Pediatr. Hematol. Oncol.* 17:425-427.
- Razin, S., and J. Tully (ed.). 1983. *Methods in Mycoplasmology*. Academic Press, New York, N.Y.
- Rechcigl, N. A., and J. E. Rechcigl. 1997. *Environmentally Safe Approaches to Crop Disease Control*. CRC Press, Boca Raton, Fla.
- Repiso, T., V. Garcia-Patos, N. Martin, M. Creus, P. Bastida, and A. Castells. 1996. Disseminated fusariosis. *Pediatr. Dermatol.* 13:118-121.
- Riley, I. T., and K. M. Ophel. 1992. *Clavibacter toxicus* sp. nov., the bacterium responsible for annual ryegrass toxicity in Australia. *Int. J. Syst. Bacteriol.* 42:64-68.
- Ritchie, B. J. 2002. Biosafety in the laboratory, p. 379-383. In J. M. Waller, J. M. Lenné, and S. J. Waller (ed.), *Plant Pathologist's Pocketbook*, 3rd ed. CABI Bioscience, Egham, United Kingdom.
- Rodriguez-Villalobos, H., M. Aoun, C. Heymans, J. M. DeBruyne, V. Duchateau, J. M. Verdebout, and F. Crokaert. 2002. Cross reaction between a pan-*Candida* genus probe and *Fusarium* spp. in a fatal case of *Fusarium oxysporum* pneumonia. *Eur. J. Clin. Microbiol. Infect. Dis.* 21:149-152.
- Rodriguez-Villalobos, H., A. Georgala, H. Beguin, C. Heymans, G. Pye, F. Crokaert, and M. Aoun. 2003. Disseminated infection due to *Cylindrocarpum* (*Fusarium*) *lichenicola* in a neutropenic patient with acute leukaemia: report of a case and review of the literature. *Eur. J. Clin. Microbiol. Infect. Dis.* 22:62-65.
- Romano, C., R. Bilenchi, C. Alessandrini, and C. Miracco. 1999. Case report: cutaneous phaeohyphomycosis caused by *Cladosporium oxysporum*. *Mycoses* 42:111-115.
- Romano, C., M. Fimiani, M. Pellegrino, L. Valenti, L. Casini, C. Miracco, and E. Faggi. 1996. Cutaneous phaeohyphomycosis due to *Alternaria tenuissima*. *Mycoses* 39:211-215.
- Romano, C., C. Miracco, and E. M. Difonzo. 1998. Skin and nail infections due to *Fusarium oxysporum* in Tuscany, Italy. *Mycoses* 41:433-437.
- Romano, C., L. Valenti, C. Miracco, C. Alessandrini, E. Paccagnini, E. Faggi, and E. M. Difonzo. 1997. Two cases of cutaneous phaeohyphomycosis by *Alternaria alternata* and *Alternaria tenuissima*. *Mycopathologia* 137:65-74.
- Rosenblueth, M., L. Martinez, J. Silva, and E. Martinez-Romero. 2004. *Klebsiella variicola*, a novel species with clinical and plant-associated isolates. *Syst. Appl. Microbiol.* 27:27-35.
- Ross, J. P., S. M. Holland, V. J. Gill, E. S. DeCarlo, and J. I. Gallin. 1995. Severe *Burkholderia* (*Pseudomonas*) *gladioli* infection in chronic granulomatous disease: report of two successfully treated cases. *Clin. Infect. Dis.* 21:1291-1293.
- Rossmann, S. N., P. L. Cernoch, and J. R. Davis. 1996. Dematiaceous fungi are an increasing cause of human disease. *Clin. Infect. Dis.* 22:73-80.
- Rott, P., and J. C. Comstock. 2000. Seedling foliage blights, p. 190-206. In P. Rott, R. Bailey, J. C. Comstock, B. Croft, and S. Saumtally (ed.), *A Guide to Sugarcane Diseases*. CIRAD Publication Service, Montpellier, France.
- Rowan, N. J., G. Caldow, C. Gemmell, and I. Hunter. 2003. Production of diarrheal enterotoxins and other potential virulence factors by veterinary isolates of *Bacillus* species associated with nongastrointestinal infections. *Appl. Environ. Microbiol.* 69:2372-2376.
- Rubinstein, I., and G. W. Pedersen. 2002. *Bacillus* species are present in chewing tobacco sold in the United States and evoke plasma exudation from the oral mucosa. *Clin. Diagn. Lab. Immunol.* 9:1057-1060.
- Saettler, A. L., N. W. Schaad, and D. A. Roth (ed.). 1989. *Detection of Bacteria in Seed and Other Planting Material*. APS Press, Inc., St. Paul, Minn.
- Saleh, O. I., P. Y. Huang, and J. S. Huang. 1997. *Bacillus pumilus*, the cause of bacterial blotch of immature Balady peach in Egypt. *J. Phytopathol.* (Berlin) 145:447-453.
- Sander, A., U. Beyer, and R. Amberg. 1998. Systemic *Fusarium oxysporum* infection in an immunocompetent patient with an adult respiratory distress syndrome (ARDS) and extracorporeal membrane oxygenation (ECMO). *Mycoses* 41:109-111.
- Satou, M., M. Ichinoe, F. Fukumoto, N. Tezuka, and S. Horiuchi. 2001. *Fusarium* blight of kangaroo paw (*Anigozanthos* spp.) caused by *Fusarium chlamydosporum* and *Fusarium semitectum*. *J. Phytopathol.* 149:203-206.
- Schaad, N. W. (ed.). 2001. *Laboratory Guide for Identification of Plant Pathogenic Bacteria*, 3rd ed. American Phytopathological Society, St. Paul, Minn.
- Schink, B., J. C. Ward, and J. G. Zeikus. 1981. Microbiology of wetwood: importance of pectin degradation and *Clostridium* species in living trees. *Appl. Environ. Microbiol.* 42:526-532.
- Schneck, N. C. (ed.). 1982. *Methods and Principles of Mycorrhizal Research*. APS Press, Inc., St. Paul, Minn.
- Scholthof, K. G. 2003. One foot in the furrow: linkages between agriculture, plant pathology, and public health. *Annu. Rev. Public Health* 24:153-174.
- Segal, B. H., T. J. Walsh, J. M. Liu, J. D. Wilson, and K. J. Kwon-Chung. 1998. Invasive infection with *Fusarium chlamydosporum* in a patient with aplastic anemia. *J. Clin. Microbiol.* 36:1772-1776.
- Seta, S., M. Gonzalez, and G. Lori. 2004. First report of walnut canker by *Fusarium incarnatum* in Argentina. *Plant Pathol.* 53:248.
- Shin, J. H., S. H. Kim, M. G. Shin, S. P. Suh, D. W. Ryang, and M. H. Jeong. 1997. Bacteremia due to *Burkholderia gladioli*: case report. *Clin. Infect. Dis.* 25:1264-1265.
- Singh, H., R. S. Singh, and J. S. Chohan. 1974. Fruit rot of *Luffa acutangula* by *Mucor circinelloides*. *Indian J. Mycol. Plant Pathol.* 4:99-100.

- Singleton, L. L., J. D. Mihail, and C. M. Rush (ed.). 1992. *Methods for Research on Soilborne Phytopathogenic Fungi*. APS Press, Inc., St. Paul, Minn.
- Smart, M. G., D. T. Wicklow, and R. W. Caldwell. 1990. Pathogenesis in *Aspergillus* ear rot of maize: light microscopy of fungal spread from wounds. *Phytopathology* 80:1287-1294.
- Sonoda, R. M. 1991. *Exserohilum rostratum* and *Bipolaris hawaiiensis* causing leaf and culm lesions on Callide Rhodesgrass. *Proc. Soil Crop Sci. Soc. Florida* 50:28-30.
- Southern, P. M., Jr. 1996. Bacteremia due to *Agrobacterium tumefaciens* (radiobacter). Report of infection in a pregnant woman and her stillborn fetus. *Diagn. Microbiol. Infect. Dis.* 24:43-45.
- Spotts, R. A., and L. A. Cervantes. 2002. Involvement of *Aureobasidium pullulans* and *Rhodotorula glutinis* in russet of d'Anjou pear fruit. *Plant Dis.* 86:625-628.
- Starr, M. P., and A. K. Chatterjee. 1972. The genus *Erwinia*: enterobacteria pathogenic to plants and animals. *Annu. Rev. Microbiol.* 26:389-426.
- Stenson, S., A. Brookner, and S. Rosenthal. 1982. Bilateral endogenous necrotizing scleritis due to *Aspergillus oryzae*. *Ann. Ophthalmol.* 14:67-72.
- St. Leger, R. J., S. E. Screen, and B. Shams-Pirzadeh. 2000. Lack of host specialization in *Aspergillus flavus*. *Appl. Environ. Microbiol.* 66:320-324.
- Su, L., J. Ou, H. Leu, P. Chiang, Y. Chiu, J. Chia, A. Kuo, C. Chiu, C. Chu, T. Wu, C. Sun, T. Riley, B. Chang, and The Infection Control Group. 2003. Extended epidemic of nosocomial urinary tract infections caused by *Serratia marcescens*. *J. Clin. Microbiol.* 41:4726-4732.
- Suckstorff, I., and G. Berg. 2003. Evidence for dose-dependent effects on plant growth by *Stenotrophomonas* strains from different origins. *J. Appl. Microbiol.* 95:656-663.
- Summerbell, R. C., S. E. Richardson, and J. Kane. 1988. *Fusarium proliferatum* as an agent of disseminated infection in an immunosuppressed patient. *J. Clin. Microbiol.* 26:82-87.
- Suominen, I., M. Andersson, A.-M. Hallaksela, and M. S. Salkinoja-Salonen. 1999. Identifying toxic *Bacillus pumilus* from industrial contaminants, food, paperboard and live trees, p. 100-105. In A. C. J. Tuijelaars, R. A. Samson, F. M. Rombouts, and S. Notermans (ed.), *Food Microbiology and Food Safety into the Next Millennium*. Proceedings of the 17th International Conference of the International Committee on Food Microbiology and Hygiene, Veldoven, The Netherlands, 13-17 September 1999.
- Taj-Aldeen, S. J., A. A. Hilal, and W. A. Schell. 2004. Allergic fungal rhinosinusitis: a report of 8 cases. *Am. J. Otolaryngol.* 25:213-218.
- Tan, M.-W. 2002. Cross-species infections and their analysis. *Annu. Rev. Microbiol.* 56:539-565.
- Tanaka, K., and F. Nonaka. 1977. Studies on the rot of onion bulbs by *Aspergillus niger*. *Proc. Assoc. Plant Prot. Kyushu* 23:36-39.
- Tang, L. M., and S. T. Chen. 1994. *Klebsiella pneumoniae* meningitis: prognostic factors. *Scand. J. Infect. Dis.* 26:95-102.
- Taylor, L. H., S. M. Latham, and M. E. Woolhouse. 2001. Risk factors for human disease emergence. *Philos. Trans. R. Soc. Lond. B.* 356:983-989.
- Tessari, G., A. Forni, R. Ferretto, M. Solbiati, G. Faggian, A. Mazzucco, and A. Barba. 2003. Lethal systemic dissemination from a cutaneous infection due to *Curvularia lunata* in a heart transplant recipient. *J. Eur. Acad. Dermatol. Venereol.* 17:440-442.
- Thind, T. S., S. B. Saksena, and S. C. Agrawal. 1976. Post harvest decay of apple fruits incited by *Aspergillus candidus* in Madhya Pradesh. *Indian Phytopathol.* 29:318.
- Thomas, P. A. 2003. Current perspectives on ophthalmic mycoses. *Clin. Microbiol. Rev.* 16:730-797.
- Thompson, J. R., K. L. Perry, and W. De Jong. 2004. A new potato virus in a new lineage of picorna-like viruses. *Arch. Virol.* 149:2141-2154.
- Toh, H., H. Hayashida, and T. Miyata. 1983. Sequence homology between retroviral reverse transcriptase and putative polymerases of hepatitis B virus and cauliflower mosaic virus. *Nature* 305:827-829.
- Torda, A. J., and P. D. Jones. 1997. Necrotizing cutaneous infection caused by *Curvularia brachyspora* in an immunocompetent host. *Australas. J. Dermatol.* 38: 85-87.
- Torii, K., Y. Noda, Y. Miyazaki, and M. Ohta. 2003. An unusual outbreak of infusion-related bacteremia in a gastrointestinal disease ward. *Jpn. J. Infect. Dis.* 56: 177-178.
- Tuite, J. 1990. Teachers supplement: Laboratory exercises of methods in plant pathology: fungi and bacteria. Purdue University, West Lafayette, Ind.
- U.S. Department of Agriculture, Animal and Plant Health Inspection Service. 2001. Federal Plant Pest Regulations. Title 7 CFR, Part 330.200. Movement of Plant Pests. http://a257.g.akamaitech.net/7/257/2422/14mar20010800/edocket.access.gpo.gov/cfr_2001/janqtr/pdf/7cfr330.100.pdf
- U.S. Department of Agriculture, Office of Agricultural Biotechnology. 1992. Guidelines for Research Involving Planned Introduction into the Environment of Genetically Modified Organisms. Guidelines recommended to USDA by the Agricultural Biotechnology Research Advisory Committee, December 3-4, 1991. <http://www.aphis.usda.gov/brs/pdf/abrac%201991.pdf>
- Usharani, P., and P. Ramarao. 1981. Corm rot of *Colocasia esculenta* caused by *Cylindrocarpon lichenicola*. *Indian Phytopathol.* 34:381-382.
- Vidaver, A. K. 1996. Emerging and re-emerging infectious diseases. *ASM News* 62:583-585.
- Vikram, H. R., E. T. Shore, and P. R. Venkatesh. 1999. Community acquired *Pseudomonas aeruginosa* pneumonia. *Conn. Med.* 63:271-273.
- Vismer, H. F., W. F. Marasas, J. P. Rheeder, and J. J. Joubert. 2002. *Fusarium dimerum* as a cause of human eye infections. *Med. Mycol.* 40:399-406.
- Watanabe, K., K. Abe, and M. Sato. 2000. Biological control of an insect pest by gut-colonizing *Enterobacter cloacae* transformed with ice nucleation gene. *J. Appl. Microbiol.* 88:90-97.
- Westcott, C. 2001. *Westcott's Plant Disease Handbook*, 6th ed. Revised by R. Kenneth Horst. Kluwer Academic Publishers, Boston, Mass.
- Wigley, P., and N. F. Burton. 1999. Genotypic and phenotypic relationships in *Burkholderia cepacia* isolated from cystic fibrosis patients and the environment. *J. Appl. Microbiol.* 86:460-468.
- Wilson, C. L., J. D. Franklin, and P. L. Pusey. 1987. Biological control of *Rhizopus* rot of peach with *Enterobacter cloacae*. *Phytopathology*. 77:303-305.
- Woods, C. W., A. M. Bressler, J. J. LiPuma, B. D. Alexander, D. A. Clements, D. J. Weber, C. M. Moore, L. B. Reller, and K. S. Kaye. 2004. Virulence associated

- with outbreak-related strains of *Burkholderia cepacia* complex among a cohort of patients with bacteremia. *Clin. Infect. Dis.* 38:1243–1250.
- Wright, K., S. Popli, V. C. Gandhi, J. R. Lentino, C. V. Reyes, and D. J. Leehey. 2003. *Paecilomyces* peritonitis: case report and review of the literature. *Clin. Nephrol.* 59:305–310.
- Yamada, K., T. Mori, S. Irie, M. Matsumura, M. Nakayama, T. Hirano, K. Suda, and K. Oshimi. 1998. Systemic aspergillosis caused by an aflatoxin-producing strain of *Aspergillus* in a post-bone marrow transplant patient with acute myeloid leukemia. *Rinsho Ketsueki* 39:1103–1108.
- Yamaguchi, M., H. Nishiya, K. Mano, O. Kunii, and H. Miyashita. 1992. Chronic necrotising pulmonary aspergillosis caused by *Aspergillus niger* in a mildly immunocompromised host. *Thorax* 47:570–571.
- Yamamoto, N., T. Matsumoto, and Y. Ishibashi. 2001. Fungal keratitis caused by *Colletotrichum gloeosporioides*. *Cornea* 20:902–903.
- Yang, S. 1973. Isolation and effect of temperature on spore germination, radial growth and pathogenicity of *Curvularia senegalensis*. *Phytopathology* 63:1540–1541.
- Yang, S. M., J. B. Morris, P. W. Unger, and T. E. Thompson. 1979. *Rhizopus* head rot of cultivated sunflower, *Helianthus annuus*, in Texas USA. *Plant Dis. Rep.* 63:833–835.
- Yeghen, T., L. Fenelon, C. K. Campbell, D. W. Warnock, A. V. Hoffbrand, H. G. Prentice, and C. C. Kibbler. 1996. *Chaetomium* pneumonia in patient with acute myeloid leukaemia. *J. Clin. Pathol.* 49:184–186.
- Yohalem, D. S., and J. W. Lorbeer. 1997. Distribution of *Burkholderia cepacia* phenotypes by niche, method of isolation and pathogenicity to onion. *Ann. Appl. Biol.* 130:467–479.
- Zinniel, D. K., P. Lambrecht, N. B. Harris, Z. Feng, D. Kuczmarski, P. Higley, C. A. Ishimaru, A. Arunakumari, R. G. Barletta, and A. K. Vidaver. 2002. Isolation and characterization of endophytic colonizing bacteria from agronomic crops and prairie plants. *Appl. Environ. Microbiol.* 68:2198–2208.